

Test Organisms	Dried Virus Control	Sample	Result	Log Reduction	CrossOver to Swine Pathogen Based on Viral Family
Avian Infectious Bronchitis virus Beaudette IB42	6.42 Log ₁₀	A	<=0.5 Log ₁₀	>=5.92 Log ₁₀	TGE and PED
		B	<=0.5 Log ₁₀	>=5.92 Log ₁₀	
	6.5 Log ₁₀	C	<=0.5 Log ₁₀	>=6.0 Log ₁₀	
Avian Influenza A (H3N2) virus (Avian Reassortant) (ATCC VR-2072)	4.75 Log ₁₀	A	<=0.5 Log ₁₀	>=4.25 Log ₁₀	Swine Influenza
		B	<=0.5 Log ₁₀	>=4.25 Log ₁₀	
		C	<=0.5 Log ₁₀	>=4.25 Log ₁₀	
Avian Influenza A (H5N1) virus	6.75 Log ₁₀	A	<=0.5 Log ₁₀	>=6.25 Log ₁₀	
		B	<=0.5 Log ₁₀	>=6.25 Log ₁₀	
Canine Coronavirus ATCC VR-809	4.5 Log ₁₀	A	<=0.5 Log ₁₀	>=4.0 Log ₁₀	TGE and PED
		B	<=0.5 Log ₁₀	>=4.0 Log ₁₀	
	4.75 Log ₁₀	C	<=0.5 Log ₁₀	>=4.25 Log ₁₀	
Canine Distemper virus	6.25 Log ₁₀	A	<=0.5 Log ₁₀	>=05.75 Log ₁₀	Nipah Virus
		B	<=0.5 Log ₁₀	>=5.75 Log ₁₀	
	6.75 Log ₁₀	C	<=0.5 Log ₁₀	>=6.25 Log ₁₀	
Feline Picornavirus	4.5 Log ₁₀	A	<=0.5 Log ₁₀	>=4.0 Log ₁₀	Swine Vesicular Exanthema
		B	<=0.5 Log ₁₀	>=4.0 Log ₁₀	
	5.75 Log ₁₀	C	<=0.5 Log ₁₀	>=5.25 Log ₁₀	
Hepatitis B Virus	5.06 Log ₁₀	A	<=0.27 Log ₁₀	>=4.79 Log ₁₀	No Crossover
	5.20 Log ₁₀	B	<=0.41 Log ₁₀	>=4.79 Log ₁₀	
	5.06 Log ₁₀	Confirmatory B	<=0.27 Log ₁₀	>=4.79 Log ₁₀	
Hepatitis C Virus	6.21 Log ₁₀	A	<=0.24 Log ₁₀	>=5.97 Log ₁₀	Classical Swine Fever
	6.21 Log ₁₀	B	<=0.42 Log ₁₀	>=5.79 Log ₁₀	
	6.06 Log ₁₀	Confirmatory B	<=0.13 Log ₁₀	>=5.93 Log ₁₀	

Xtreme raw material Base was evaluated in the presence of 5% serum and 400 ppm hard water with 10 minute contact time and found to be effective against the above noted viruses on hard, nonporous environmental surfaces.

Herpes Simplex Virus Type 1	5.5 Log ₁₀	A	<=0.5 Log ₁₀	>=5.0 Log ₁₀	No Crossover
		B	<=0.5 Log ₁₀	>=5.0 Log ₁₀	
	6.0 Log ₁₀	C	<=0.5 Log ₁₀	>=5.5 Log ₁₀	

Xtreme raw material Base was evaluated in the presence of 5% serum and 400 ppm hard water with 10 minute contact time and found to be effective against the above noted viruses on hard, nonporous environmental surfaces.

Herpes Simplex Virus Type 2	6.0 Log ₁₀	A	<=0.5 Log ₁₀	>=5.5 Log ₁₀	No Crossover
		B	<=0.5 Log ₁₀	>=5.5 Log ₁₀	
	5.75 Log ₁₀	C	<=0.5 Log ₁₀	>=5.25 Log ₁₀	
Human Coronavirus	4.5 Log ₁₀	A	<=0.5 Log ₁₀	>=4.0 Log ₁₀	No Crossover
		B	<=0.5 Log ₁₀	>=4.0 Log ₁₀	
	4.5 Log ₁₀	C	<=0.5 Log ₁₀	>=4.0 Log ₁₀	
Human Immunodeficiency Virus type 1 (HIV 1)	5.75 Log ₁₀	A	<=1.5 Log ₁₀	>=4.25 Log ₁₀	No Crossover
		B	<=1.5 Log ₁₀	>=4.25 Log ₁₀	
		C	<=1.5 Log ₁₀	>=4.25 Log ₁₀	
Infectious Bovine Rhinotracheitis virus	4.5 Log ₁₀	A	<=0.0 Log ₁₀	>=4.0 Log ₁₀	See Pseudorabies
		B	<=0.0 Log ₁₀	>=4.0 Log ₁₀	
	4.75 Log ₁₀	C	<=0.0 Log ₁₀	>=4.25 Log ₁₀	
Influenza A virus	6.5 Log ₁₀	A	<=0.0 Log ₁₀	>=6.0 Log ₁₀	Swine Influenza
		B	<=0.0 Log ₁₀	>=6.0 Log ₁₀	
	6.0 Log ₁₀	C	<=0.0 Log ₁₀	>=5.5 Log ₁₀	
Pseudorabies virus	6.25 Log ₁₀	A	<=0.5 Log ₁₀	>=5.75 Log ₁₀	Swine Pathogen
		B	<=0.5 Log ₁₀	>=5.75 Log ₁₀	
	5.5 Log ₁₀	C	<=0.5 Log ₁₀	>=5.0 Log ₁₀	
Respiratory Syncytial virus ATCC VR-26	4.5 Log ₁₀	A	<=0.5 Log ₁₀	>=4.0 Log ₁₀	Blue eye disease
		B	<=0.5 Log ₁₀	>=4.0 Log ₁₀	
	5.0 Log ₁₀	C	<=0.5 Log ₁₀	>=4.5 Log ₁₀	
Transmissible Gastroenteritis virus	4.75 Log ₁₀	A	<=0.5 Log ₁₀	>=4.35 Log ₁₀	Swine Pathogen
		B	<=0.5 Log ₁₀	>=4.25 Log ₁₀	
	6.25 Log ₁₀	C	<=0.5 Log ₁₀	>=5.75 Log ₁₀	
Vaccinia virus	6.75 Log ₁₀	A	<=0.5 Log ₁₀	>=6.25 Log ₁₀	Swine Pox
		B	<=0.5 Log ₁₀	>=6.25 Log ₁₀	
	6.5 Log ₁₀	C	<=0.5 Log ₁₀	>=6.0 Log ₁₀	

Xtreme raw material Base was evaluated in the presence of 5% serum and 400 ppm hard water with 10 minute contact time and found to be effective against the above noted viruses on hard, nonporous environmental surfaces.

Test Organisms	Dried Virus Control	Sample	Result	Log Reduction	
Canine Parvovirus Type 2b	7.5 Log ₁₀	A	<=3.5 Log ₁₀	>=4.0 Log ₁₀	Swine Parvovirus
		B	<=3.5 Log ₁₀	>=4.0 Log ₁₀	

Xtreme raw material Base was evaluated in the presence of 5% serum and 400 ppm hard water with 10 minute contact time and found to be effective against the above noted viruses on hard, nonporous environmental surfaces.